

Important Information - Please Read Before Using This Policy

The following coverage policy applies to health benefit plans administered by PHP and may not be covered by all PHP plans. Please refer to the member's benefit document for specific coverage information. If there is a difference between this general information and the member's benefit document, the member's benefit document will be used to determine coverage. For example, a member's benefit document may contain a specific exclusion related to a topic addressed in a coverage policy.

Coverage determinations for individual requests require consideration of:

- The terms of the applicable benefit document in effect on the date of service.
- Any applicable laws and regulations.
- Any relevant collateral source materials including coverage policies.
- The specific facts of the particular situation.

Contact PHP Customer Service to discuss plan benefits more specifically.

1.0 Policy:

Health Plan covers preventive health services to adults and children, as required by the ACA.

Certain preventive health services require prior approval for coverage.

2.0 Background

The Affordable Care Act (ACA) requires individual and group health plans to cover network preventive health services and immunizations without a member cost-share (e.g., deductibles, coinsurance, and copayments) unless the plan qualifies under the grandfather provision or for an exemption. Coverage for preventive health services other than those mandated by ACA is dependent on benefit plan language.

The preventive health services listed in this policy are based on recommendations from:

- United States Preventive Task Force (USPSTF), grade A or B recommendations.
- Centers for Disease Control and Prevention (CDC).

- Health Resources and Services Administration (HRSA).
- Latest medical research from organizations like the American Medical Association (AMA).
- Pharmacy Benefit Manager.

3.0 Clinical Determination Guidelines:

- 1. Preventive health services do not incur a member cost share when they are all the following:
 - a. Listed in this policy.
 - b. Done for preventive purposes (not diagnostic).
 - c. Completed by a network provider.
- 2. Some services MAY require prior approval. See table below.
 - a. Screening versus diagnostic, monitoring or surveillance testing.

A positive result on a preventive screening exam does not alter its classification as a preventive service but does influence how that service is classified when rendered in the future. For example, if a screening colonoscopy performed on an asymptomatic individual without additional risk factors for colorectal cancer (e.g., adenomatous polyps, inflammatory bowel disease) detects colorectal cancer or polyps, the purpose of the procedure remains screening, even if polyps are removed during the preventive screening. However, once a diagnosis of colorectal cancer or additional risk factors for colorectal cancer is identified, future colonoscopies will no longer be considered preventive screening.

3. Reporting of preventive care services.

Preventive health services are reported with diagnosis and procedure codes, which identify the services as preventive and not for treatment of injury or illness (see code chart below). Age, gender or frequency limits are utilized for certain designated services (i.e., wellness exams, vision and hearing screening, administration of Human Papillomavirus [HPV] vaccines, nutritional and genetic counseling). Preventive health services submitted with diagnosis codes that represent treatment of illness or injury are paid as applicable under member's normal medical benefits rather than preventive care coverage.

4.0 Coding:

Prior Approval Legend: Y = All lines of business; N = None required; 1 = HMO/POS; 2 = EPO/PPO; 3 = SHS 264; 4 = SHS 1269 Non-Union & Union; 5 = LBWL; 6 = Dart; 7 = SHS 1269 Union Only.

Medical Services

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|--|--------------|--|----------------|-----------------------|----------------|---|--------|------------------------------------|-------|
| Abdominal Aortic Aneurysm (AAA) Screening | 76706 | Ultrasound, abdominal aorta, real time with image documentation; screening study for abdominal aortic aneurysm (AAA) | | х | F17.200 | Nicotine dependence, unspecified, uncomplicated | х | | |
| (12/20/19 USPSTF | | | | | F17.201 | Nicotine dependence, unspecified, in remission | | | |
| recommenda- tion for men aged 65 – 75 who have ever | | | | | F17.210 | Nicotine dependence, cigarettes, uncomplicated | | | |
| smoked, one- time screening) | | | | | F17.211 | Nicotine dependence, cigarettes, in remission | | | |
| | | | | | F17.220 | Nicotine dependence, chewing tobacco, uncomplicated | | | |
| | | | | | F17.221 | Nicotine dependence, chewing tobacco, in remission | | | |
| | | | | | F17.290 | Nicotine dependence, other tobacco product, uncomplicated | | | |
| | | | | | F17.291 | Nicotine dependence, other tobacco product, in remission | | | |
| | | | | | Z87.891 | Personal history of nicotine dependence | | | |
| Alcohol/ Substance Misuse Screening and Behavioral | 99408 | Alcohol and/or Substance Abuse structured screening, and brief intervention services; 15 to 30 minutes | | Х | Z72.89 | Other problems related to lifestyle | Х | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|---|----------------|-----------------------|----------------|---|--------|------------------------------------|-------|
| Counseling | 99409 | Alcohol and/or Substance Abuse structured screening, and brief intervention services; >30 minutes | | х | Z000.0 | Encounter for general adult medical examination | х | | |
| | G0396 | Alcohol and/or substance abuse structured assessment, and brief intervention 15 to 30 minutes | | х | Z000.1 | Encounter for general adult medical examination with abnormal findings | x | | |
| | G0397 | Alcohol and/or substance abuse structured assessment, and brief intervention > 30 minutes | | х | Z013.89 | Encounter for screening for other disorder | х | | |
| | G0442 | Annual alcohol misuse screening, 15 minutes | | х | | | х | | |
| | G0443 | Brief face-to-face behavioral counseling for alcohol misuse, 15 minutes | | x | | | х | | |
| Annual physical | 96160 | Administration of patient- focused health risk assessment instrument (e.g., health hazard appraisal) with scoring and documentation, per standardized instrument | | | | | х | | |
| | 96161 | Administration of caregiver- focused health risk assessment instrument (e.g., depression inventory) for the benefit of the patient, with scoring and documentation, per standardized instrument | | | | | х | | |
| | 99381 | Initial Preventive Medicine E&M, new patient; Infant (younger than 1 year) | | | | | | х | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|---|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | 99382 | Initial Preventive Medicine E&M, new patient; Early childhood (1-4 yrs) | | | | | | х | |
| | 99383 | Initial Preventive Medicine E&M, new patient; Late childhood (5-11 yrs) | | | | | | х | |
| | 99384 | Initial Preventive Medicine E&M, new patient; Adolescent (12-17 yrs) | | | | | | х | |
| | 99385 | Initial Preventive Medicine E&M, new patient; 18-39 yrs | | | | | x | | |
| | 99386 | Initial Preventive Medicine E&M, new patient; 40-64 yrs | | | | | Х | | |
| | 99387 | Initial Preventive Medicine E&M, new patient; 65 yrs and older | | | | | x | | |
| | 99391 | Periodic Preventive Medicine E&M, established patient; Infant (younger than 1 year) | | | | | | х | |
| | 99392 | Periodic Preventive Medicine E&M, established patient; Early childhood (1-4 yrs) | | | | | | x | |
| | 99393 | Periodic Preventive Med E&M, established patient; Late childhood (5-11 yrs) | | | | | | х | |
| | 99394 | Periodic Preventive Medicine E&M, established patient; Adolescent (12-17 yrs) | | | | | | x | |
| | 99395 | Periodic Preventive Medicine E&M, established patient; 18- 39 yrs | | | | | x | | |
| | 99396 | Periodic Preventive Medicine E&M, established patient; 40- 64 yrs | | | | | x | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|---|--------------|---|----------------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | 99397 | Periodic Preventive Medicine E&M, established patient; 65 yrs and older | | | | | x | | |
| | 99401 | Preventive Medicine Counseling, individual; 15 minutes | | | | | х | | |
| | 99402 | Preventive Medicine Counseling, individual; 30 minutes | | | | | х | | |
| | 99403 | Preventive Medicine Counseling, individual; 45 minutes | | | | | x | | |
| | 99404 | Preventive Medicine Counseling, individual; 60 minutes | | | | | х | | |
| | 99411 | Preventive Medicine Counseling, group; 30 minutes | | | | | x | | |
| | 99412 | Preventive Medicine Counseling, group; 60 minutes | | | | | х | | |
| | 99429 | Unlisted preventive medicine service | Y - MRM to review | | | | x | | |
| | 99461 | Initial care, E&M normal newborn infant seen in other than hospital or birthing center | | | | | | х | |
| | G0468 | Federally qualified health center visit, initial preventive physical exam or annual wellness visit | | | | | x | | |
| Cardio-vascular screening (includes blood | 80061 | Lipid Panel | | х | Z13.6 | Encounter for screening for cardiovascular disorders | х | | |
| pressure and | 82465 | Cholesterol | | Х | | | Х | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------------------|--------------|--|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| labs) | 84478 | Triglycerides | | Х | | | Х | | |
| Colorectal cancer screening | 00811 | Anesthesia for lower intestinal endoscopic procedures, endoscope introduced distal to duodenum; not otherwise specified | | x | Z00.00 | Encounter for general adult medical examination without abnormal findings | x | | |
| | 00812 | Anesthesia for lower intestinal endoscopic procedures, endoscope introduced distal to duodenum; screening colonoscopy | | х | Z00.01 | Encounter for general adult medical examination with abnormal findings | х | | |
| | 44392 | Colonoscopy-Stoma; w/ removal of tumor, polyp, lesion, forceps | | х | Z12.10 | Encounter for screening for malignant neoplasm of intestinal tract, unspecified | x | | |
| | 44394 | Colonoscopy-Stoma; w/ removal of tumor, polyp, lesion, snare | | х | Z12.11 | Encounter for screening for malignant neoplasm of colon | x | | |
| | 44401 | Colonoscopy-stoma; w/ ablation of tumor(s), polyp(s), or other lesion(s) | | x | Z12.12 | Encounter for screening for malignant neoplasm of rectum | х | | |
| | 45300 | Proctosigmoidoscopy, rigid; diagnostic | | х | Z12.13 | Encounter for screening for malignant neoplasm of small intestine | х | | |
| | 45305 | Proctosigmoidoscopy, rigid; diagnostic, with biopsy, single or multiple | | х | Z80.0 | Family history of malignant neoplasm of digestive organs | х | | |
| | 45308 | Proctosigmoidoscopy, rigid; diagnostic, w/ removal of single tumor, polyp, or lesion, forceps or bipolar cautery | | х | Z83.71 | Family history of colonic polyps | х | | |
| | 45309 | Proctosigmoidoscopy, rigid; diagnostic, with removal of single tumor, polyp, lesion, snare | | х | Z85.00 | Personal history of malignant neoplasm of unspecified digestive organ | х | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|--|----------------|-----------------------|----------------|---|--------|------------------------------------|-------|
| | 45315 | Proctosigmoidoscopy, rigid; diagnostic, w/ removal of multiple tumors, polyps, lesions, forceps, snare | | х | Z85.038 | Personal history of other malignant neoplasm of large intestine | х | | |
| | 45320 | Proctosigmoidoscopy, rigid; diagnostic, w/ ablation of tumor, polyp, lesion by forceps, bipolar cautery or snare | | х | Z85.048 | Personal history of malignant neoplasm of epididymis | х | | |
| | 45330 | Sigmoidoscopy flexible; diagnostic | | х | Z86.010 | Personal history of colonic polyps | Х | | |
| | 45331 | Sigmoidoscopy flexible; diagnostic w/ biopsy, single or multiple | | х | | | х | | |
| | 45332 | Sigmoidoscopy flexible; diagnostic w/ removal foreign body | | х | | | х | | |
| | 45333 | Sigmoidoscopy flexible; diagnostic w/ removal of tumor, polyp, lesion by forceps | | х | | | х | | |
| | 45334 | Sigmoidoscopy flexible; diagnostic w/ control of bleeding, any method | | х | | | х | | |
| | 45338 | Sigmoidoscopy flexible; diagnostic w/ removal of tumor, polyp, lesion by snare | | x | | | х | | |
| | 45346 | Sigmoidoscopy, flexible; with ablation of tumor(s), polyp(s), or other lesion(s) | | х | | | х | | |
| | 45341 | Sigmoidoscopy flexible; diagnostic w/ endoscopic ultrasound | | х | | | х | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|---|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | 45342 | Sigmoidoscopy flexible; diagnostic w/ transendoscopic ultrasound guided fine needle aspiration/biopsy | | х | | | x | | |
| | 45378 | Colonoscopy flexible; diagnostic including collection of specimen by brushing or washing | | х | | | x | | |
| | 45380 | Colonoscopy flexible; diagnostic w/ biopsy, single or multiple | | х | | | x | | |
| | 45381 | Colonoscopy flexible; diagnostic w/ directed submucosal injection(s) | | x | | | x | | |
| | 45382 | Colonoscopy flexible; diagnostic w/ control of bleeding, any method | | x | | | x | | |
| | 45388 | Colonoscopy flexible; diagnostic w/ ablation of tumor, polyp, lesion | | x | | | x | | |
| | 45384 | Colonoscopy flexible; diagnostic w/ removal of tumor, polyp, lesion by forceps | | x | | | x | | |
| | 45385 | Colonoscopy flexible; diagnostic w/ removal of tumor, polyp, lesion by snare | | x | | | x | | |
| | 81528 | Oncology (colorectal) screening, quantitative real- time target and signal amplification of 10 DNA markers (KRAS mutations, promoter methylation of NDRG4 and BMP3) and fecal hemoglobin, utilizing stool, algorithm report as a positive | | Х | | | x | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|---|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | | or negative result (i.e., Cologuard) | | | | | | | |
| | 82270 | Blood occult, by peroxidase activity, feces | | х | | | Х | | |
| | G0104 | Colorectal cancer screening; flexible sigmoidoscopy | | | | | Х | | |
| | G0105 | Colorectal cancer screening; colonoscopy, high risk | | | | | Х | | |
| | G0106 | Colorectal cancer screening; alternative to G0104, screening sigmoidoscopy, barium enema | | | | | х | | |
| | G0120 | Colorectal cancer screening; alternative to G0105, screening colonoscopy, barium enema | | | | | х | | |
| | G0121 | Colorectal cancer screening; colonoscopy on individual not meeting criteria for high risk | | | | | х | | |
| | G0122 | Colorectal cancer screening; barium enema | | | | | Х | | |
| | G0328 | Colorectal cancer screening; fecal-occult blood test, immunoassay, 1-3 simultaneous | | | | | х | | |
| Contraception | 58300 | Insertion of intrauterine device (IUD) | | | Z30013 | Encounter for initial prescription of injectable contraceptive | | | х |
| | 58301 | Removal of intrauterine device (IUD) | | x | Z30019 | Encounter for initial prescription of contraceptives, unspecified | | | х |
| | 11981 | Insertion non-biodegradable drug delivery implant | | | Z3009 | Encounter for other general counseling and advice on contraception | | | х |
| | 57170 | Diaphragm or cervical cap | | | Z3040 | Encounter for | | | Х |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-------------------------|--------------|---|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | | fitting w/ instructions | | | | surveillance of contraceptives, unspecified | | | |
| | 96372 | Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); subcutaneous or intramuscular | | x | Z3042 | Encounter for surveillance of injectable contraceptive | | | х |
| | A4261 | Cervical cap for contraceptive use | | | Z308 | Encounter for other contraceptive management | | | Х |
| | A4264 | Permanent implantable contraceptive intratubal occlusion device(s) and delivery system | | | Z309 | Encounter for contraceptive management, unspecified | | | Х |
| | 58300 | Insertion of intrauterine device (IUD) | | | Z30013 | Encounter for initial prescription of injectable contraceptive | | | Х |
| | S4981 | Insertion of levonorgestrel- releasing intrauterine system | | | | | | | Х |
| | S4989 | Contraceptive intrauterine device, including implants and supplies | | | | | | | Х |
| Depression Screening | 96127 | Brief emotional/ behavioral assessment, with scoring and documentation, per standardized instrument | | х | Z13.89 | Encounter for screening for other disorder | х | х | |
| | G0444 | Annual depression screening 15 minutes | | х | | | х | х | |
| Diabetes Screening | 82947 | Glucose; quantitative, blood (except reagent strip) | | х | Z13.1 | Encounter for screening for diabetes mellitus | х | | |
| | 82948 | Glucose; blood, reagent strip | | Х | | | Х | | |
| | 82950 | Glucose; post glucose dose (includes glucose) | | Х | | | х | | |
| | 82951 | Glucose; tolerance test 3 specimens | | х | | | х | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|---|--------------|--|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | 83036 | Hemoglobin; glycated (A1C) | | Х | | | Х | | |
| | 0403T | Preventive behavior changes, intensive program of prevention of diabetes using a standardized diabetes prevention program curriculum, provided to individuals in a group setting, minimum 60 minutes, per day | | | | | x | | |
| | 0488T | Preventive behavior changes, online/ electronic structured intensive program for prevention of diabetes using a standardized diabetes prevention program curriculum, provided to an individual, per 30 days | | | | | x | | |
| Immunizations (Vaccines) | G0008 | Administration of influenza virus vaccine | | | | | x | | |
| | G0009 | Administration of pneumococcal vaccine | | | | | X | Х | |
| | G0010 | Administration of Hepatitis B vaccine | | | | | x | Х | |
| Infectious and sexually transmitted | 99401 | Preventive Medicine Counseling, individual, 15 minutes | | | | | x | | |
| disease counseling | 99402 | Preventive Medicine Counseling, individual, 30 minutes | | | | | x | | |
| | 99403 | Preventive Medicine Counseling, individual, 45 minutes | | | | | x | | |
| | 99404 | Preventive Medicine Counseling, individual, 60 minutes | | | | | x | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|--|--------------|---|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | 99411 | Preventive Medicine Counseling, group, 30 minutes | | | | | x | | |
| | 99412 | Preventive Medicine Counseling, group, 60 minutes | | | | | x | | |
| | G0445 | High intensity behavioral counseling to prevent sexually transmitted infection, individual, semi-annual, 30 minutes | | | | | x | | |
| Infectious and sexually transmitted disease | 86631 | Antibody; Chlamydia | | x | Z00.00 | Encounter for general adult medical examination w/o abnormal findings | x | | |
| screening: Chlamydia | 86632 | Antibody; Chlamydia, IgM | | х | Z00.01 | Encounter for general adult medical examination w/ abnormal findings | x | | |
| | 87110 | Culture, chlamydia, any source | | х | Z00.8 | Encounter for other general examination | х | | |
| | 87270 | Infectious agent antigen detection by immunofluorescent technique; Chlamydia trachomatis | | × | Z01.411 | Encounter for gynecological examination (general) (routine) w/ abnormal findings | x | | |
| | 87320 | Infectious agent antigen detection by immunoassay technique, (e.g., enzyme immunoassay [EIA], enzyme- linked immunosorbent assay [ELISA], immunochemiluminometric assay [IMCA]) qualitative or semiquantitative, multiple- step method; Chlamydia trachomatis | | Х | Z01.419 | Encounter for gynecological examination (routine) w/o abnormal findings | x | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|--|--------------|--|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | 87490 | Infectious agent detection by nucleic acid (DNA or RNA); Chlamydia trachomatis, direct probe technique | | х | Z01.42 | Encounter for cervical smear to confirm findings of recent normal smear following initial abnormal smear | x | | |
| | 87491 | Infectious agent detection by nucleic acid (DNA or RNA); Chlamydia trachomatis, amplified probe technique | | х | Z32.00 | Encounter for pregnancy test, result unknown | х | | |
| | 87492 | Infectious agent detection by nucleic acid (DNA or RNA); Chlamydia trachomatis, quantification | | х | Z32.01 | Encounter for pregnancy test, result positive | х | | |
| | 87810 | Infectious agent antigen detection by immunoassay with direct optical observation; Chlamydia trachomatis | | х | Z32.02 | Encounter for pregnancy test, result negative | х | | |
| | | | | | Z11.8 | Encounter for screening for other infectious and parasitic diseases | | | |
| | | | | | Z11.59 | Encounter for screening for other viral diseases | | | |
| | | | | | Z11.3 | Encounter for screening for infections with a predominantly sexual mode of transmission | | | |
| Infectious and sexually transmitted disease | 87590 | Infectious agent detection by nucleic acid (DNA or RNA); Neisseria gonorrhoeae, direct probe technique | | х | Z00.00 | Encounter for general adult medical examination w/o abnormal findings | х | | |
| screening: Gonorrhea | 87591 | Infectious agent detection by nucleic acid (DNA or RNA); Neisseria gonorrhoeae, amplified probe technique | | х | Z00.01 | Encounter for general adult medical examination w/ abnormal findings | Х | | |
| | 87592 | Infectious agent detection by nucleic acid (DNA or RNA); Neisseria gonorrhoeae, | | х | Z00.8 | Encounter for other general examination | Х | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|--|--------------|--|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | | quantification | | | | | | | |
| | 87850 | Infectious agent antigen detection by immunoassay with direct optical observation; Neisseria gonorrhoeae | | x | Z01.411 | Encounter for gynecological examination (routine) w/ abnormal findings | X | | |
| | | | | | Z01.419 | Encounter for gynecological examination (routine) w/o abnormal findings | | | |
| | | | | | Z01.42 | Encounter for cervical smear to confirm findings of recent normal smear following initial abnormal smear | | | |
| | | | | | Z11.3 | Encounter for screening for infections with a predominantly sexual mode of transmission | | | |
| Infectious and sexually transmitted disease | 86704 | Hepatitis B core antibody (HBcAb); total | | x | Z00.00 | Encounter for general adult medical examination w/o abnormal findings | х | | |
| screening: Hepatitis B | 86705 | Hepatitis B core antibody (HBcAb); IgM antibody | | x | Z00.01 | Encounter for general adult medical examination w/ abnormal findings | х | | |
| | 86706 | Hepatitis B surface antibody (HBsAb) | | x | Z00.8 | Encounter for other general examination | х | | |
| | 86707 | Hepatitis Be antibody (HBeAb) | | x | Z01.411 | Encounter for gynecological examination (routine) w/ abnormal findings | х | | |
| | 87340 | Infectious agent antigen detection by immunoassay technique, (e.g., enzyme immunoassay [EIA], enzyme- | | x | Z01.419 | Encounter for gynecological examination (routine) w/o abnormal findings | х | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|--|--------------|---|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | | linked immunosorbent assay [ELISA], immunochemiluminometric assay [IMCA]) qualitative or semiquantitative, multiple- step method; hepatitis B surface antigen (HBsAg) | | | | | | | |
| | 87341 | Infectious agent antigen detection by immunoassay technique, (e.g., enzyme immunoassay [EIA], enzyme- linked immunosorbent assay [ELISA], immunochemiluminometric assay [IMCA]) qualitative or semiquantitative, multiple- step method; hepatitis B surface antigen (HBsAg) neutralization | | x | Z11.3 | Encounter for screening for infections with a predominantly sexual mode of transmission | x | | |
| | 87516 | Infectious agent detection by nucleic acid (DNA or RNA); hepatitis B virus, amplified probe technique | | x | Z77.21 | Contact w/ and (suspected) exposure to potentially hazardous body fluids | х | | |
| | 87517 | Infectious agent detection by nucleic acid (DNA or RNA); hepatitis B virus, quantification | | х | | | х | | |
| | G0499 | Hepatitis B screening in non- pregnant, high risk individual | | х | | | х | | |
| Infectious and sexually transmitted disease | 86803 | Hepatitis C antibody | | x | Z00.00 | Encounter for general adult medical examination w/o abnormal findings | x | | |
| screening: Hepatitis C | 86804 | Hepatitis C antibody; confirmatory test (e.g., immunoblot) | | х | Z00.01 | Encounter for general adult medical examination w/ abnormal findings | х | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|--|--------------|--|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | 87520 | Infectious agent detection by nucleic acid (DNA or RNA); hepatitis C, direct probe technique | | х | Z00.8 | Encounter for other general examination | х | | |
| | 87521 | Infectious agent detection by nucleic acid (DNA or RNA); hepatitis C, amplified probe technique, includes reverse transcription when performed | | х | Z01.411 | Encounter for gynecological examination (routine) w/ abnormal findings | х | | |
| | 87522 | Infectious agent detection by nucleic acid (DNA or RNA); hepatitis C, quantification, includes reverse transcription when performed | | х | Z01.419 | Encounter for gynecological examination (routine) w/o abnormal findings | x | | |
| | 87902 | Infectious agent genotype analysis by nucleic acid (DNA or RNA); Hepatitis C virus | | Х | Z11.3 | Encounter for screening for infections with a predominantly sexual mode of transmission | Х | | |
| | G0472 | Hepatitis C antibody screening for individual at high risk and other covered indication(s) | | Х | Z77.21 | Contact w/ and (suspected) exposure to potentially hazardous body fluids | Х | | |
| Infectious and sexually transmitted disease | 86689 | Antibody; HTLV or HIV antibody, confirmatory test (e.g., Western Blot | | x | Z00.00 | Encounter for general adult medical examination w/o abnormal findings | x | | |
| screening: HIV/ AIDS | 86701 | Antibody; HIV-1 | | х | Z00.01 | Encounter for general adult medical examination w/ abnormal findings | х | | |
| | 86702 | Antibody; HIV-2 | | х | Z00.8 | Encounter for other general examination | х | | |
| | 86703 | Antibody; HIV-1 and HIV-2, single result | | х | Z01.411 | Encounter for gynecological examination (routine) w/ abnormal findings | х | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|--|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | 87389 | Infectious agent antigen detection by immunoassay technique, (e.g., enzyme immunoassay [EIA], enzyme- linked immunosorbent assay [ELISA], immunochemiluminometric assay [IMCA]) qualitative or semiquantitative, multiple- step method; HIV-1 antigen(s), with HIV-1 and HIV-2 antibodies, single result | | x | Z01.419 | Encounter for gynecological examination (routine) w/o abnormal findings | x | | |
| | 87390 | Infectious agent antigen detection by immunoassay technique, (e.g., enzyme immunoassay [EIA], enzyme- linked immunosorbent assay [ELISA], immunochemiluminometric assay [IMCA]) qualitative or semiquantitative, multiple- step method; HIV-1 | | x | Z11.3 | Encounter for screening for infections with a predominantly sexual mode of transmission | x | | |
| | 87391 | Infectious agent antigen detection by immunoassay technique, (e.g., enzyme immunoassay [EIA], enzyme- linked immunosorbent assay [ELISA], immunochemiluminometric assay [IMCA]) qualitative or semiquantitative, multiple- step method; HIV-2 | | X | | | x | | |
| | 87534 | Infectious agent detection by nucleic acid (DNA or RNA); HIV-1, direct probe technique | | х | | | х | | |
| | 87535 | Infectious agent detection by nucleic acid (DNA or RNA); HIV-1, amplified probe | | х | | | х | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|--|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | | technique, includes reverse transcription when performed | | | | | | | |
| | 87536 | Infectious agent detection by nucleic acid (DNA or RNA); HIV-1, quantification, includes reverse transcription when performed | | x | | | x | | |
| | 87537 | Infectious agent detection by nucleic acid (DNA or RNA); HIV-2, direct probe technique | | х | | | х | | |
| | 87538 | Infectious agent detection by nucleic acid (DNA or RNA); HIV-2, amplified probe technique, includes reverse transcription when performed | | x | | | x | | |
| | 87539 | Infectious agent detection by nucleic acid (DNA or RNA); HIV-2, quantification, includes reverse transcription when performed | | х | | | x | | |
| | G0432 | Infectious agent antibody detection by enzyme immunoassay (EIA) technique, HIV-1 and/or HIV- 2, screening | | х | | | x | | |
| | G0433 | Infectious agent antibody detection by enzyme-linked immunosorbent assay (ELISA) technique, HIV-1 and/or HIV-2, screening | | | | | x | | |
| | G0435 | Infectious agent antigen detection by rapid antibody test of oral mucosa transudate, HIV-1 or HIV-2, screening | | х | | | x | | |
| | G0475 | HIV antigen/antibody, combination assay, screening | | x | | | х | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|--|--------------|--|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| Infectious and sexually transmitted disease screening: | 87623 | Infectious agent detection by nucleic acid (DNA or RNA); Human Papillomavirus (HPV), low-risk types (e.g., 6, 11, 42, 43, 44) | | x | Z00.00 | Encounter for general adult medical examination w/o abnormal findings | x | | |
| Human papillomavirus (HPV) | 87624 | Infectious agent detection by nucleic acid (DNA or RNA); Human Papillomavirus (HPV), high-risk types (e.g., 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 68) | | Х | Z00.01 | Encounter for general adult medical examination w/ abnormal findings | х | | |
| | 87625 | Infectious agent detection by nucleic acid (DNA or RNA); Human Papillomavirus (HPV), types 16 and 18 only, includes type 45, if performed | | x | Z00.8 | Encounter for other general examination | x | | |
| | | | | | Z01.411 | Encounter for gynecological examination (routine) w/ abnormal findings | | | |
| | | | | | Z01.419 | Encounter for gynecological examination (routine) w/o abnormal findings | | | |
| | | | | | Z11.3 | Encounter for screening for infections with a predominantly sexual mode of transmission | | | |
| | | | | | Z11.51 | Encounter for screening for human papillomavirus (HPV) | | | |
| | | | | | Z12.4 | Encounter for screening for malignant neoplasm of cervix | | | |
| | | | | | Z12.72 | Encounter for screening for malignant neoplasm of vagina | | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|--|--------------|---|----------------|-----------------------|----------------------|--|--------|------------------------------------|-------|
| Infectious and sexually transmitted disease | 86592 | Syphilis test, non-treponemal antibody; qualitative (e.g., VDRL, RPR, ART) | | х | Z00.00 | Encounter for general adult medical examination w/o abnormal findings | х | | |
| screening: Syphilis | 86593 | Syphilis test, non-treponemal antibody; quantitative | | х | Z00.01 | Encounter for general adult medical examination w/ abnormal findings | Х | | |
| | 86780 | Antibody; Treponema pallidum | | х | Z00.8 | Encounter for other general examination | Х | | |
| | | | | | Z01.411 | Encounter for gynecological examination (routine) w/ abnormal findings | | | |
| | | | | | Z01.419 | Encounter for gynecological examination (routine) w/o abnormal findings | | | |
| | | | | | Z11.3 | Encounter for screening for infections with a predominantly sexual mode of transmission | | | |
| | | | | | Z32.00 | Encounter for pregnancy test, result unknown | | | |
| | | | | | Z32.01 | Encounter for pregnancy test, result positive | | | |
| | | | | | Z32.02 | Encounter for pregnancy test, result negative | | | |
| Lung cancer screening | 71250 | CT; thorax, w/o contrast material | | | F17.200 - F17.299 | Nicotine Dependence | Х | | |
| | G0296 | Counseling visit to discuss need for lung cancer screening using low dose CT scan (LDCT) (service is for eligibility determination and shared decision making) | | | Z12.2 | Encounter for screening for malignant neoplasm of respiratory organs | х | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|--|--------------|---|----------------|-----------------------|-------------------|---|--------|------------------------------------|-------|
| | G0297 | Low dose CT scan for lung cancer screening | | | Z87.891 | Personal history of nicotine dependence | | | |
| Nutritional counseling | G0446 | Intensive behavioral therapy for cardiovascular disease, individual, annual, 15 minutes | | x | Z00.00 | Encounter for general adult medical exam without abnormal findings | x | x | |
| | G0447 | Behavioral counseling for obesity, 15 minutes | | x | Z00.01 | Encounter for general adult medical exam with abnormal findings | x | x | |
| | G0473 | Face-to-face behavioral counseling for obesity, group (2-10), 30 minutes | | х | Z00.121 | Encounter for routine child health exam with abnormal findings | х | х | |
| | | | | | Z00.129 | Encounter for routine child health exam without abnormal findings | | | |
| | | | | | Z00.8 | Encounter for other general exam | | | |
| | | | | | Z68.2 – Z68.54 | Body mass index (BMI) | | | |
| | | | | | Z71.3 | Dietary counseling and surveillance | | | |
| Obesity screening and counseling | 99401 | Preventive Medicine Counseling, individual; 15 minutes | | х | Z13.89 | Encounter for screening for other disorder | х | Х | |
| | 99402 | Preventive Medicine Counseling, individual; 30 minutes | | х | Z68.30 | Body mass index (BMI) 30.0 – 30.9, adult | x | х | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|---|--------------|---|----------------|-----------------------|----------------|---|--------|------------------------------------|-------|
| | 99403 | Preventive Medicine Counseling, individual; 45 minutes | | x | Z68.39 | Body mass index (BMI) 39.0 – 39.9, adult | x | x | |
| | 99404 | Preventive Medicine Counseling, individual; 60 minutes | | х | Z68.41 | Body mass index (BMI) 40.0 – 44.9, adult | х | х | |
| | 99411 | Preventive Medicine Counseling, group; 30 minutes | | х | Z68.45 | Body mass index (BMI) 70 or greater, adult | х | х | |
| | 99412 | Preventive Medicine Counseling, group; 60 minutes | | х | | | x | х | |
| | G0447 | Behavioral counseling for obesity, 15 minutes | | х | | | х | х | |
| Prostate cancer screening | 84152 | Prostate specific antigen (PSA); complexed (direct measurement) | | | | | х | | |
| | 84153 | Prostate specific antigen (PSA); total | | | | | Х | | |
| | 84154 | Prostate specific antigen (PSA); free | | | | | Х | | |
| | G0102 | Prostate cancer screening; digital rectal exam | | | | | х | | |
| | G0103 | Prostate cancer screening; PSA test | | | | | х | | |
| Skin cancer behavioral counseling | 99401 | Preventive Medicine Counseling, individual; 15 minutes | | | | | x | x | |
| | 99402 | Preventive Medicine Counseling, individual; 30 minutes | | | | | x | х | |
| | 99403 | Preventive Medicine Counseling, individual; 45 minutes | | | | | х | х | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|------------------------------|--------------|--|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | 99404 | Preventive Medicine Counseling, individual; 60 minutes | | | | | x | х | |
| | 99411 | Preventive Medicine Counseling, group; 30 minutes | | | | | х | х | |
| | 99412 | Preventive Medicine Counseling, group; 60 minutes | | | | | х | х | |
| Tobacco use counseling | 99406 | Smoking and tobacco use cessation counseling visit; intermediate, 3 to 10 minutes | | | | | x | | |
| | 99407 | Smoking and tobacco use cessation counseling visit; intermediate, greater than 10 minutes | | | | | x | | |
| Tuberculosis (TB) testing | 86480 | Tuberculosis test | | x | Z00.00 | Encounter for general adult medical examination w/o abnormal findings | х | | |
| | 86580 | Skin test; tuberculosis intradermal | | х | Z00.01 | Encounter for general adult medical examination w/ abnormal findings | Х | | |
| | | | | | Z03.89 | Encounter for observation for other suspected diseases and conditions ruled out | | | |
| | | | | | Z11.1 | Encounter for screening for respiratory tuberculosis | | | |
| Developmental screening | 96110 | Developmental screening, with scoring and documentation, per standardized instrument | | х | Z00.110 | Health examination for newborn under 8 days old | | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|-------------------------------------|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | | | | | Z00.111 | Health examination for newborn 8 to 28 days old | | | |
| | | | | | Z00.121 | Encounter for routine child health examination with abnormal findings | | | |
| | | | | | Z00.129 | Encounter for routine child health examination without abnormal findings | | | |
| | | | | | Z13.40 | Encounter for screening for unspecified developmental delays | | | |
| | | | | | Z13.41 | Encounter for autism screening | | | |
| | | | | | Z13.42 | Encounter for screening for global developmental delays (milestones) | | | |
| | | | | | Z13.49 | Encounter for screening for other developmental delays | | | |
| | | | | | Z76.2 | Encounter for health supervision and care of other healthy infant and child | | | |
| Hearing screening | 92551 | Screening test, pure tone, air only | | x | Z00.110 | Health examination for newborn under 8 days old | | х | |
| | V5008 | Hearing screening | | x | Z00.111 | Health examination for newborn 8 to 28 days old | | х | |
| | | | | | Z00.121 | Encounter for routine child health examination with abnormal findings | | х | |
| | | | | | Z00.129 | Encounter for routine child health examination without abnormal findings | | Х | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------------------|--------------|--|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | | | | | Z76.2 | Encounter for health supervision and care of other healthy infant and child | | х | |
| Hypo- thyroidism Screening | 84436 | Thyroxine; total | | x | Z13.29 | Encounter for screening for other suspected endocrine disorder | | х | |
| | 84437 | Thyroxine; requiring elution (e.g., neonatal) | | | | | | Х | |
| | 84439 | Thyroxine; free | | | | | | Х | |
| | 84443 | Thyroid Stimulating Hormone | | | | | | Х | |
| Lead screening | 83655 | Lead | | x | Z13.88 | Encounter for screening for disorder due to exposure to contaminants | | x | |
| Newborn metabolic screening | S3620 | Newborn metabolic screening panel, includes test kit, postage and lab tests specified by the state for inclusion in this panel | | x | Z00.110 | Health examination for newborn under 8 days old | | x | |
| | | | | | Z00.111 | Health examination for newborn 8 to 28 days old | | | |
| Oral Health | 99188 | Application of topical fluoride varnish by a physician or other qualified health care professional | | | | | | х | |
| PKU screening | 83498 | Hydroxyprogesterone 17-D | | x | Z13.228 | Encounter for screening for other metabolic disorders | | x | |
| | 84030 | Phenylalanine (PKU), blood | | Х | | | | Х | |
| Sickle cell screening | 83020 | Hemoglobin fractionation and quantitation; electrophoresis (e.g., A2, S, C, and/or F) | | x | Z13.0 | Encounter for screening for diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism | | х | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|---------------------------------|--------------|---|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | 83021 | Hemoglobin fractionation and quantitation; chromatography (e.g., A2, S, C, and/or F) | | x | | | | х | |
| | S3850 | Genetic testing for sickle cell anemia | | х | | | | х | |
| Vision screening | 99172 | Visual function screening | | x | Z00.110 | Health examination for newborn under 8 days old | | х | |
| | 99173 | Screening test of visual acuity | | x | Z00.111 | Health examination for newborn 8 to 28 days old | | х | |
| | G0117 | Glaucoma screening for high risk patients furnished by an optometrist or ophthalmologist | | x | Z00.121 | Encounter for routine child health examination with abnormal findings | | х | |
| | G0118 | Glaucoma screening for high risk patients furnished under the direct supervision of an optometrist or ophthalmologist | | × | Z00.129 | Encounter for routine child health examination without abnormal findings | | х | |
| | | | | | Z76.2 | Encounter for health supervision and care of other healthy infant and child | | | |
| Asymptomatic Bacteriuria in | 81007 | Urinalysis; bacteriuria screen | | х | O09.00 | Supervision of high risk pregnancy | | | Х |
| Pregnant Women, Screening | 87081 | Culture, presumptive, pathogenic organisms, screening only | | x | O09.93 | Supervision of high risk pregnancy | | | Х |
| | 87084 | Culture, presumptive, pathogenic organisms, screening only; with colony estimation from density chart | | x | Z33.1 | Pregnant state, incidental | | | х |
| | 87086 | Culture, bacterial; quantitative colony count, urine | | x | Z34.00 | Encounter for supervision of normal first pregnancy, unspecified trimester | | | х |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|--|--------------|--|----------------|-----------------------|----------------|--|--------|------------------------------------|-------|
| | 87088 | Culture, bacterial; with isolation and presumptive identification of each isolate, urine | | х | Z34.01- | Encounter for supervision of normal first pregnancy, first trimester | | | Х |
| | | | | | Z34.93 | Encounter for supervision of normal pregnancy, unspecified, third trimester | | | |
| Breast Cancer Screening: BRCA Genetic Counseling | 96040 | Medical Genetics and Genetic Counseling Services, each 30 minutes, face-to-face with patient/family | | | | | | | Х |
| | 99401 | Preventive Medicine Counseling, individual; 15 minutes | | | | | | | Х |
| | 99402 | Preventive Medicine Counseling, individual; 30 minutes | | | | | | | Х |
| | 99403 | Preventive Medicine Counseling, individual; 45 minutes | | | | | | | х |
| | 99404 | Preventive Medicine Counseling, individual; 60 minutes | | | | | | | х |
| Breast Cancer Screening: BRCA Genetic Mutation Testing | 81211 | BRCA1, BRCA2 gene analysis; full sequence analysis and common duplication/deletion variants in BRCA 1 | х | | | | | | х |
| | 81163 | BRCA1 (BRCA1, DNA repair associated), BRCA2 (BRCA2, DNA repair associated) (eg, hereditary breast and ovarian cancer) gene analysis; full sequence analysis | х | | | | | | Х |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|---|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | 81164 | BRCA1 (BRCA1, DNA repair associated), BRCA2 (BRCA2, DNA repair associated) (eg, hereditary breast and ovarian cancer) gene analysis; full duplication/deletion analysis (ie, detection of large gene rearrangements) | Х | | | | | | х |
| | 81165 | BRCA1 (BRCA1, DNA repair associated) (eg, hereditary breast and ovarian cancer) gene analysis; full sequence analysis | Х | | | | | | х |
| | 81166 | BRCA1 (BRCA1, DNA repair associated) (eg, hereditary breast and ovarian cancer) gene analysis; full duplication/deletion analysis (ie, detection of large gene rearrangements) | х | | | | | | x |
| | 81167 | BRCA2 (BRCA2, DNA repair associated) (eg, hereditary breast and ovarian cancer) gene analysis; full duplication/deletion analysis (ie, detection of large gene rearrangements) | Х | | | | | | х |
| | 81212 | BRCA1, BRCA2 (e.g., hereditary breast and ovarian cancer) gene analysis; 185delAG, 5385insC, 6174del variants | х | | | | | | х |
| | 81215 | BRCA1 (breast cancer 1) (e.g., hereditary breast and ovarian cancer) gene analysis; known familial variant | х | | | | | | x |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|--|--------------|---|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | 81216 | BRCA2 (breast cancer 2) (e.g., hereditary breast and ovarian cancer); full sequence analysis | х | | | | | | х |
| | 81217 | BRCA2 (breast cancer 2) (e.g., hereditary breast and ovarian cancer); known familial variant | х | | | | | | Х |
| | 81162 | BRCA1, BRCA2 (breast cancer 1 and 2) (e.g., hereditary breast and ovarian cancer) gene analysis; full sequence analysis and full duplication/deletion analysis | Х | | | | | | х |
| Breast Cancer Screening: Mammography | 77067 | Screening mammography, bilateral, including computer- aided detection (CAD) | | | | | | | х |
| Breast Cancer Screening: Risk Reduction of Primary Breast Cancer | | Tamoxifen or Raloxifene | х | | | | | | х |
| Domestic Violence Screening/ | 99401 | Preventive Medicine Counseling, individual; 15 | | | | | | | х |
| counseling | 99402 | Preventive Medicine Counseling, individual; 30 minutes | | | | | | | х |
| | 99403 | Preventive Medicine Counseling, individual; 45 minutes | | | | | | | Х |
| | 99404 | Preventive Medicine Counseling, individual; 60 minutes | | | | | | | х |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|--|--------------|--|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | 99411 | Preventive Medicine Counseling, group; 30 minutes | | | | | | | Х |
| | 99412 | Preventive Medicine Counseling, group; 60 minutes | | | | | | | х |
| Gynecological Exam: Cervical Cancer Screening | 88141 | Cytopathology, cervical or vaginal (any reporting system), requiring interpretation by physician | | | | | | | х |
| | 88142 | Cytopathology, cervical or vaginal (any reporting system), collected in preservative fluid, automated thin layer preparation; manual screening under physician supervision | | | | | | | x |
| | 88143 | Cytopathology, cervical or vaginal (any reporting system), collected in preservative fluid, automated thin layer preparation; with manual screening and rescreening under physician supervision | | | | | | | x |
| | 88147 | Cytopathology smears, cervical or vaginal; screening by automated system under physician supervision | | | | | | | х |
| | 88148 | Cytopathology smears, cervical or vaginal; screening by automated system with manual rescreening under physician supervision | | | | | | | х |
| | 88150 | Cytopathology, slides, cervical or vaginal; manual screening under physician supervision | | | | | | | х |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|---|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | 88152 | Cytopathology, slides, cervical or vaginal; with manual screening and computer-assisted rescreening under physician supervision | | | | | | | х |
| | 88153 | Cytopathology, slides, cervical or vaginal; with manual screening and rescreening under physician supervision | | | | | | | х |
| | 88155 | Cytopathology, slides, cervical or vaginal, definitive hormonal evaluation (e.g., maturation index, karyopyknotic index, estrogenic index) | | | | | | | x |
| | 88164 | Cytopathology, slides, cervical or vaginal (the Bethesda System); manual screening under physician supervision | | | | | | | х |
| | 88165 | Cytopathology, slides, cervical or vaginal (the Bethesda System); with manual screening and rescreening under physician supervision | | | | | | | х |
| | 88166 | Cytopathology, slides, cervical or vaginal (the Bethesda System); with manual screening and computer-assisted rescreening under physician supervision | | | | | | | x |
| | 88167 | Cytopathology, slides, cervical or vaginal (the Bethesda System); with | | | | | | | Х |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|--|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | | manual screening and computer-assisted rescreening using cell selection and review under physician supervision | | | | | | | |
| | 88174 | Cytopathology, cervical or vaginal (any reporting system), collected in preservative fluid, automated thin layer preparation; screening by automated system, under physician supervision | | | | | | | x |
| | 88175 | Cytopathology, cervical or vaginal (any reporting system), collected in preservative fluid, automated thin layer preparation; with screening by automated system and manual rescreening or review, under physician supervision | | | | | | | x |
| | G0101 | Cervical or vaginal cancer screening; pelvic and clinical breast exam | | | | | | | х |
| | G0123 | Screening cytopathology | | | | | | | Х |
| | G0124 | cervical or vaginal | | | | | | | Х |
| | G0141 | Screening cytopathology smears, cervical or vaginal, performed by automated system, with manual rescreening, requiring interpretation by physician | | | | | | | х |
| | G0143 | Screening cytopathology, cervical or vaginal (any reporting system), collected in preservative fluid, automated thin layer | | | | | | | х |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|---|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | | preparation, with manual screening and rescreening by cytotechnologist under physician supervision | | | | | | | |
| | G0144 | Screening cytopathology, cervical or vaginal (any reporting system), collected in preservative fluid, automated thin layer preparation, with screening by automated system, under physician supervision | | | | | | | Х |
| | G0145 | Screening cytopathology, cervical or vaginal (any reporting system), collected in preservative fluid, automated thin layer preparation, with screening by automated system and manual rescreening under physician supervision | | | | | | | х |
| | G0147 | Screening cytopathology smears, cervical or vaginal, performed by automated system under physician supervision | | | | | | | х |
| | G0148 | Screening cytopathology smears, cervical or vaginal, performed by automated system with manual rescreening | | | | | | | х |
| | P3000 | Screening Pap smear, cervical or vaginal | | | | | | | Х |
| | P3001 | Screening Papanicolaou smear, cervical or vaginal, up to 3 smears, requiring interpretation by physician | | | | | | | х |
| | Q0091 | Screening Papanicolaou smear; obtaining, preparing | | | | | | | Х |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|---|--------------|--|----------------|-----------------------|--------------------|--|--------|------------------------------------|-------|
| | | and conveyance of cervical or vaginal smear to laboratory | | | | | | | |
| | S0610 | Annual gynecological exam, new patient | | | | | | | Х |
| | S0612 | Annual gynecological exam, established patient | | | | | | | Х |
| | S0613 | Annual gynecological exam; clinical breast exam without pelvic exam | | | | | | | х |
| Osteoporosis Screening: Bone Density Study | 76977 | Ultrasound bone density measurement & interpretation, peripheral site(s), any method | | x | Z13.820 | Encounter for screening for osteoporosis | | | Х |
| | 77078 | Computed tomography (CT), bone mineral density study, 1 or more sites; axial skeleton | | x | | | | | х |
| | 77080 | Dual-Energy X-Ray Absorptiometry (DXA), Bone Density Study, 1 or More Sites; Axial Skeleton | | x | | | | | х |
| | 77085 | Dual-energy X-ray absorptiometry (DXA), bone density study, 1 or more sites | | x | | | | | Х |
| | G0130 | Single energy x-ray absorptiometry bone density study, one or more sites | | | | | | | х |
| Pregnancy: Anemia screening | 85013 | Blood count, spun microhematocrit | | х | Z13.0 | Encounter for screening for diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism | | | х |
| | 85014 | Blood count; hematocrit (Hct) | | x | Z34.00 - Z34.93 | Encounter for supervision of normal pregnancy | | | Х |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|---|--------------|--|----------------|-----------------------|--------------------------------------|---|--------|------------------------------------|-------|
| | 85018 | Blood count; hemoglobin (Hgb) | | | | | | | Х |
| Pregnancy: Breastfeeding counseling | S9443 | Lactation class non-physician provider, per session | | | | | | | х |
| Pregnancy: Breast Pump | E0602 | Breast pump, manual, any type | | | | | | | х |
| | E0603 | Breast pump, electric, any type | | | | | | | х |
| | E0604 | Breast pump, hospital grade, electric, any type | | | | | | | х |
| Pregnancy: Labs for Pre- | 82947 | Glucose; quantitative, blood (except reagent strip) | | x | O09.00 - O09.03 | Supervision of high risk pregnancy | | | х |
| and Post-natal Care and | 82948 | Glucose; blood, reagent strip | | х | O30.00 - O30.93 | Multiple gestation | | | х |
| Delivery | 82950 | Glucose; post glucose dose (includes glucose) | | x | Z34.00 - Z34.93 | Encounter for supervision of normal pregnancy | | | х |
| | 82951 | Glucose; tolerance test (GTT), 3 specimens (includes glucose) | | x | Z36.0- Z36.5, Z36.81- Z36.9 | Encounter for antenatal screening | | | х |
| | 80055 | Obstetric panel | | х | Z37.0 - Z37.9 | Outcome of delivery | | | х |
| | 80081 | Obstetric panel (includes HIV testing) | | x | Z39.0 - Z39.2 | Encounter for maternal postpartum care and exam | | | х |
| | 86901 | Blood typing; RH | | Х | | | | | Х |
| Female Sterilization Procedures | 58565 | Hysteroscopy, surgical; with bilateral fallopian tube cannulation to induce occlusion by placement of permanent implants | | | | | | | х |
| | 58600 | Ligation or transection of fallopian tube(s), abdominal | | | | | | | Х |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|--|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | | or vaginal approach, unilateral or bilateral | | | | | | | |
| | 58605 | Ligation or transection of fallopian tube(s), abdominal or vaginal approach, postpartum, unilateral or bilateral, during same hospitalization | | | | | | | х |
| | 58611 | Ligation or transection of fallopian tube(s) when done at the time of cesarean delivery or intra-abdominal surgery (not a separate procedure) | | | | | | | х |
| | 58615 | Occlusion of fallopian tube(s) by device (e.g., band, clip, Falope ring) vaginal or suprapubic approach | | | | | | | х |
| | 58670 | Laparoscopy, surgical; with fulguration of oviducts (with or without transection) | | | | | | | Х |
| | 58671 | Laparoscopy, surgical; with occlusion of oviducts by device | | | | | | | х |

Prescription Drugs

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|---|--------------|--|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| Cardio- vascular; blood clot/stroke prevention for men and women ages 50-59 | | Generic over the counter Aspirin 81mg with a once daily dosing | | | | | х | | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|--|--------------|--|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | | Atorvastatin 10 mg and 20 mg | | | | | х | | |
| Low to | | Fluvastatin 20 mg and 40 mg | | | | | Х | | |
| moderate dose statins for adult | | Fluvastatin ER 80 mg | | | | | Х | | |
| men and women age 40 | | Lovastatin 10 mg, 20 mg and 40 mg | | | | | х | | |
| to 75 with one or more cardiovascular | | Pravastatin 10 mg, 20 mg, 40 mg and 80 mg | | | | | х | | |
| disease (CVD) risk factors | | Rosuvastatin 5 mg and 10 mg | | | | | Х | | |
| ISK TACIOIS | | Simvastatin 5 mg, 10 mg, 20 mg and 40 mg | | | | | x | | |
| Bowel Prep for men and women ages 50 through 74 | | Citrate of Magnesia ClearLax Disposable enema Docusate Calcium GaviLyte-C GaviLyte-H and Bisacodyl GaviLyte-N Magnesium Hydroxide | | | | | x | | |
| Smoking cessation; quantity limit: 168-day supply of each product annually, *must be 18 or older | | Chantix *Nicotine Patch – generic only *Nicotine Gum – generic only *Nicotine Lozenges – generic only Bupropion SR – generic for Zyban only | | | | | x | | |
| Oral Health for children less than 5 years of age | | Generic prescription single entity or combo products providing 0.5mg per day of Fluoride | | | | | | х | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|---|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | 99188 | Application of topical fluoride varnish by a physician or other qualified health care professional | | | | | | x | |
| Contraception | J1050 | Medroxyprogesterone acetate, 1 mg | | | | | | | х |
| | J7296 | Levonorgestrel-releasing intrauterine contraceptive system (Kyleena), 19.5 mg | | | | | | | х |
| | J7297 | Levonorgestrel-releasing intrauterine contraceptive system (Liletta), 18.6 mg | | | | | | | х |
| | J7300 | Intrauterine copper contraceptive | | | | | | | х |
| | J7301 | Levonorgestrel-releasing intrauterine contraceptive system (Skyla), 13.5 mg | | | | | | | Х |
| | J7302 | Levonorgestrel-releasing intrauterine contraceptive system (Mirena), 52 mg | | | | | | | х |
| | J7303 | Contraceptive supply, hormone containing vaginal ring, each | | | | | | | х |
| | J7304 | Contraceptive supply, hormone containing patch, each | | | | | | | х |
| | J7306 | Levonorgestrel (contraceptive) implant system, including implants and supplies | | | | | | | x |
| | J7307 | Etonogestrel (contraceptive) implant system, including implant and supplies | | | | | | | х |
| Contraception | | Apri | | | | | | | Х |
| Prescription | | Camila | | | | | | | Х |
| Medications | | Conceptrol | | | | | | | Х |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|---|--------------|--|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | | Enpresse | | | | | | | Х |
| | | FC2 female condom | | | | | | | Х |
| | | Gynol II | | | | | | | Х |
| | | Junel Fe | | | | | | | Х |
| | | Kariva | | | | | | | Х |
| | | Levonorgestrel | | | | | | | Х |
| | | Low-Ogestrel | | | | | | | Х |
| | | Natazia | | | | | | | Х |
| | | NuvaRing | | | | | | | Х |
| | | Sprintec | | | | | | | Х |
| | | Today Sponge | | | | | | | Х |
| | | Tri-Sprintec | | | | | | | Х |
| | | Vaginal contraceptive film/ foam (VCF) | | | | | | | Х |
| | | Velivet | | | | | | | Х |
| | | Xulane patch | | | | | | | Х |
| Bone Health for women more than 65 years of age | | Vitamin D2 or D3 single content of 1000IU per dose or dual combination agents containing calcium and vitamin D2 or D3. | | | | | | | х |
| Prevention of Morbidity/ Mortality from Pre-eclampsia for women ages 12-59 | | Aspirin 81mg | | | | | | | x |
| Pregnancy: | | Total B With C | | | | | | | Х |
| Pre-natal | | Foltabs 800 | | | | | | | Х |
| Vitamins for | | Folic acid | | | | | | | Х |
| women less | | Dialyvite 800 | | | | | | | Х |
| han 55 years | | Stress Formula | | | | | | | Х |
| of age Immunizations | | Prenatal Complete | | | | | | | Х |
| (Vaccines) | | Perry Prenatal | | | | | | | Х |
| (1.400,100) | | Daily Prenatal | | | | | | | Х |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|--|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | | Prenatal | | | | | | | Х |
| | | KPN Prenatal | | | | | | | Х |
| | | Prenavite | | | | | | | Х |
| | | B-Balanced | | | | | | | Х |
| | | B Complex | | | | | | | Х |
| | | Vitamin B Complex | | | | | | | Х |
| Vaccines | 90460 | Immunization administration through 18 years of age via any route of administration, with counseling by physician or other qualified health care professional; first or only component of each vaccine or toxoid administered | | | | | x | х | |
| | 90461 | Immunization administration through 18 years of age via any route of administration, with counseling by physician or other qualified health care professional; each additional vaccine or toxoid component administered | | | | | x | х | |
| | 90471 | Immunization administration (includes percutaneous, intradermal, subcutaneous, or intramuscular injections); 1 vaccine (single or combination vaccine/toxoid) | | | | | X | х | |
| | 90472 | Immunization administration (includes percutaneous, intradermal, subcutaneous, or intramuscular injections); each additional vaccine | | | | | x | х | |
| | 90473 | Immunization administration by intranasal or oral route; 1 vaccine (single or combination vaccine/toxoid) | | | | | x | х | |
| | 90474 | Immunization administration | | | | | Х | Х | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|--|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | | by intranasal or oral route; each additional vaccine (single or combination vaccine/toxoid) | | | | | | | |
| | 90620 | Meningococcal recombinant protein and outer membrane vesicle vaccine, serogroup B (MenB-4C), 2 dose schedule, for intramuscular use (Bexsero – only ages 10-25 yrs.) | | | | | x | Х | |
| | 90621 | Meningococcal recombinant lipoprotein vaccine, serogroup B (MenB-FHbp), 2 or 3 dose schedule, for intramuscular use (Trumenba only for ages 10-25 yrs.) | | | | | x | x | |
| | 90630 | Influenza virus vaccine, quadrivalent, split virus, preservative free, intradermal use; 18-64 y.o. | | | | | x | | |
| | 90632 | Hepatitis A vaccine (HepA), adult dosage, IM | | | | | Х | | |
| | 90633 | Hepatitis A vaccine (HepA), pediatric/adolescent dosage- 2 dose schedule, IM | | | | | | х | |
| | 90634 | Hepatitis A vaccine (HepA), pediatric/adolescent dosage- 3 dose schedule, IM | | | | | х | х | |
| | 90636 | HepA & HepB vaccine adult dose, IM | | | | | Х | | |
| | 90644 | Meningococcal conjugate vaccine, and Haemophilus influenzae type b (Hib) vaccine, 4 dose schedule, children 6 weeks-18 months of age, IM | | | | | x | х | |
| | 90647 | Haemophilus influenzae type b vaccine (Hib), 3 dose | | | | | Х | х | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|--|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | | schedule, IM | | | | | | | |
| | 90648 | Haemophilus influenzae type b vaccine (Hib), 4 dose schedule, IM | | | | | х | х | |
| | 90649 | Human Papillomavirus (HPV) vaccine, 3 dose schedule, IM, ages 9-45 | | | | | х | х | |
| | 90650 | Human Papillomavirus (HPV) vaccine, 3 dose schedule, IM, ages 9-45 | | | | | х | х | |
| | 90651 | Human Papillomavirus (HPV) vaccine, 3 dose schedule, IM, ages 9-45 | | | | | х | х | |
| | 90653 | Influenza vaccine, IM (65 & older) | | | | | Х | | |
| | 90654 | Influenza virus vaccine, intradermal use | | | | | | х | |
| | 90655 | Influenza virus vaccine, 0.25 ml, IM | | | | | Х | х | |
| | 90656 | Influenza virus vaccine 0.5 ml, IM | | | | | | Х | |
| | 90657 | Influenza virus vaccine, children 0.25 ml, IM | | | | | Х | х | |
| | 90658 | Influenza virus vaccine, 0.5 ml, IM | | | | | Х | х | |
| | 90660 | Influenza virus vaccine, intranasal use | | | | | X | x | |
| | 90661 | Influenza virus vaccine, 0.5 ml, IM | | | | | Х | X | |
| | 90662 | Influenza virus vaccine, IM | | | | | Х | Х | |
| | 90670 | Pneumococcal conjugate vaccine, IM | | | | | Х | X | |
| | 90672 | Influenza virus vaccine, intranasal use | | | | | Х | X | |
| | 90673 | Influenza virus vaccine, IM | | | | | Х | Х | |
| | 90674 | Influenza virus vaccine, quadrivalent (ccIIV4), derived from cell cultures, subunit, | | | | | Х | х | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|--|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | | preservative and antibiotic free, 0.5 mL dosage, for intramuscular use (Flucelvax) | | | | | | | |
| | 90680 | Rotavirus vaccine, 3 dose schedule, oral use | | | | | Х | Х | |
| | 90681 | Rotavirus vaccine, oral use | | | | | Х | Х | |
| | 90682 | Influenza virus vaccine, quadrivalent (RIV4), derived from recombinant DNA, hemagglutinin (HA) protein only, preservative and antibiotic free, for intramuscular use (Flublok) | | | | | x | x | |
| | 90685 | Influenza virus vaccine, 0.25 ml, IM | | | | | | Х | |
| | 90686 | Influenza virus vaccine, quadrivalent (IIV4), split virus, preservative free, 0.5 mL dosage, for intramuscular use | | | | | | x | |
| | 90687 | Influenza virus vaccine, children 6-35 months of age, IM | | | | | | x | |
| | 90688 | Influenza virus vaccine, 0.5 ml, IM | | | | | | х | |
| | 90696 | Diphtheria, tetanus toxoids, acellular pertussis (DTaP) vaccine and inactivated poliovirus vaccine (IPV); children 4 through 6 years of age, IM | | | | | x | х | |
| | 90697 | DTaP-IPV-Hib-HepB vaccine, IM | | | | | Х | Х | |
| | 90698 | DTaP-IPV, Hib, IM | | | | | | Х | |
| | 90700 | DTaP vaccine, younger than 7 years, IM | | | | | | х | |
| | 90702 | Diphtheria and tetanus toxoids (DT) when administered to individuals | | | | | х | Х | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|---|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | | younger than 7 years, IM | | | | | | | |
| | 90707 | Measles, mumps & rubella (MMR), subq | | | | | Х | X | |
| | 90710 | Measles, mumps, rubella & varicella (MMRV), subq | | | | | Х | X | |
| | 90713 | Poliovirus vaccine, Subq | | | | | Х | Х | |
| | 90714 | Tetanus and diphtheria toxoids (Td) 7 years or older, IM | | | | | х | x | |
| | 90715 | Tetanus, diphtheria toxoids and acellular pertussis vaccine (TDaP), 7 years or older, IM | | | | | x | x | |
| | 90716 | Varicella virus vaccine (VAR), subq | | | | | Х | | |
| | 90723 | DTaP-HepB-IPV), IM | | | | | Х | Х | |
| | 90732 | Pneumococcal vaccine, 2 years or older, subq or IM | | | | | Х | x | |
| | 90733 | Meningococcal vaccine, subq | | | | | Х | Х | |
| | 90734 | Meningococcal vaccine, IM | | | | | Х | Х | |
| | 90736 | Zoster (shingles) vaccine, subq, covered for ages 50 and older | | | | | x | | |
| | 90739 | Hepatitis B vaccine (HepB), IM | | | | | Х | | |
| | 90740 | Hepatitis B vaccine (HepB), dialysis or immunosuppressed patient, IM | | | | | x | x | |
| | 90743 | Hepatitis B vaccine (HepB), adolescent, IM | | | | | Х | х | |
| | 90744 | Hepatitis B vaccine (HepB), pediatric/adolescent dosage, IM | | | | | Х | x | |
| | 90746 | Hepatitis B vaccine (HepB), adult dosage, IM | | | | | Х | | |
| | 90747 | Hepatitis B vaccine (HepB), dialysis or | | | | | Х | X | |

| Preventive Service | CPT/ HCPC | CPT/HCPC/Drug Description | Requires PA | Requires Diagnosis | ICD-10 Code | ICD-10 Description | Adults | Children (newborn to 18 yrs) | Women |
|-----------------------|--------------|--|----------------|-----------------------|----------------|--------------------|--------|------------------------------------|-------|
| | | immunosuppressed patient dosage, IM | | | | | | | |
| | 90748 | Hepatitis B and Haemophilus influenzae type b vaccine (Hib-HepB), IM | | | | | Х | х | |
| | 90750 | Zoster vaccine recombinant, adjuvanted, suspension IM (Shingrix), covered for ages 50 and older | | | | | х | | |
| | 90756 | Influenza virus vaccine, quadrivalent (ccIIV4) | | | | | Х | Х | |

5.0 Unique Configuration/Prior Approval/Coverage Details:

Preventive health claims require the modifier 33 to be considered payable as a preventive service. However, PHP does not process claims based solely on the presence of modifier 33.

Preventive health services are dependent upon claim submission using preventive diagnosis (when applicable) and procedure codes in order to be identified and covered as preventive health services.

G0438 – Annual wellness visit; includes a personalized prevention plan of service, initial visit covered as a Preventive Service for Metal Plan members.

G0439 – Annual wellness visit, includes a personalized prevention plan of service, subsequent visit covered as a Preventive Service for Metal Plan members.

6.0 Terms & Definitions:

<u>Diagnostic service</u>. Done to monitor, diagnose or treat a health problem. A deductible, copayment or coinsurance may apply. The following are examples of diagnostic services:

- Management of a chronic condition such as diabetes.
- Follow-up tests ordered by a doctor after a preventive screening determined a health problem.
- Follow-up tests ordered by a doctor based on symptoms, such as abdominal pain.

<u>Preventive health service</u>. Screenings, tests, and services performed for symptom-free or disease-free individuals. They may also include immunizations and screening services for individuals who are symptom-free or disease-free and are at increased risk for a particular disease. There is no cost to the member.

Wellness examination. Well-baby, well-child, well-adult (including well-woman) examinations that include:

- An age- and gender-appropriate history.
- Physical examinations.
- Counseling/anticipatory guidance.
- Risk factor reduction interventions.
- The ordering of appropriate immunizations and laboratory/screening procedures.

7.0 References, Citations & Resources:

- 1. HeathCare.gov Preventive health services. Available at: https://www.healthcare.gov/coverage/preventive-care-benefits/.
- 2. National Conference of State Legislatures, Preventive Services Covered Under the Affordable Care Act. Available at: <u>http://www.ncsl.org/research/health/american-health-benefit-exchanges-b.aspx</u>.
- 3. U.S. Preventive Services Task Force A and B Recommendations, available at: <u>https://www.uspreventiveservicestaskforce.org/Page/Name/uspstf-a-and-b-recommendations/</u>.
- 4. Pharmacy uses Lexicomp and Facts and Comparison.[®]
- 5. Pharmacy Breast Cancer prophylactic policy.

8.0 Associated Documents [For internal use only]:

Sample Letter – UM Approval or Denial letter for services requiring prior approval.

Forms – MMF-04 Medical Prior Authorization Form, MMF-05 Pharmacy Authorization Form.

9.0 Revision History

Original Effective Date: January 1, 2016

Next Revision Date: 01/01/2021

| Revision Date | Reason for Revision |
|---------------|--|
| 7/17 | Updated with new or more comprehensive coding and descriptions. |
| 1/18 | CPT code added for Cologuard (81528). AMA code changes effective 1/1/18: 1 code description change (90686), 4 codes deleted (G0202 - see 77067, Q9984 - see J7296, 87515 - no replcmt, 88154 - no replcmt). 3 new codes added (0403T, 0488T, 00812). Removed diagnosis requirement for 0403T and 0488T. Added new HCPC code J7296 for Kyleena. Added new CPT code 90756 Influenza virus vaccine, quadrivalent (cclIV4), 0.5mL dosage, for intramuscular Effective 1/1/2018 the following medications will be available to members with a ACA plan for 40-70 years of age for \$0 copay: |

| Revision Date | Reason for Revision |
|----------------------|---|
| | Atorvastatin 10 mg and 20 mg |
| | Fluvastatin 20 mg and 40 mg |
| | Fluvastatin ER 80 mg |
| | Lovastatin 10 mg, 20 mg and 40 mg |
| | Pravastatin 10 mg, 20 mg, 40 mg and 80 mg |
| | Rosuvastatin 5 mg and 10 mg Simvastatin 5 mg, 10 mg, 20 mg and 40 mg |
| | NOTE: For members through 39 years of age and members age 71 and greater, the copay still applies at the Tier 1 benefit. It was a mandate by the ACA that we have medication in this category covered at no cost to the member. This is the standard CVS list which was approved at the 12/6/17 Pharmacy & Therapeutics committee. |
| 6/18 | Added code 90750 for shingles vaccine. |
| 8/18 | Removed nutritional therapy codes; 97802, 97803, 97804, S9449, S9452, S9470. Added G0473. Annual review by QI/MRM 12/12/18; added immunization codes: 90620, 90621, 90674 and 90682; added anesthesia for colonoscopy code 00811. |
| 1/2019 | 1/1/2019 new codes added for BRCA: 81163-81167, strikethrough: 81211, 81213, and 81214. Added ICD-10 diagnosis codes for pre- and post-natal lab testing and for billing with G0446. |
| 2/2019 | Updated age on iron supplements and answered vaccine question |
| 11/19 | Annual review; separated medical and drugs into 2 tables, removed deleted codes from 2018 and 2019, revised age limits for HPV vaccine. |